

The Smallest Drops of the Hottest Matter
Exploring the Small Size Limit of the Quark Gluon Plasma

Anne M. Sickles, University of Illinois Urbana-Champaign

Sambamurti Memorial Lecture, July 15, 2014

how hot?

how hot?

80 ° F

how hot?

80 ° F

2000 ° F

how hot?

80 ° F

2000 ° F



28M ° F

how hot?

80 ° F

2000 ° F



28M ° F



150B ° F

how hot?

80 ° F

2000 ° F



28M ° F



150B ° F

quark-gluon plasma

no photo available

5T ° F

The nucleus



- >99% of the mass of atoms, and thus normal matter, is in the nucleus
- composed of protons and neutrons, *nucleons*
- these nucleons are held together by the strong force
  - one of the 4 fundamental forces of nature
  - characterized by very strong short range interactions

and what's inside protons and neutrons?

fundamental particles which interact
via the strong force

quarks and gluons



valence quarks define the hadron
also a **sea** of (anti)quarks & gluons

and what's inside protons and neutrons?

fundamental particles which interact
via the strong force

quarks and gluons

confinement

quarks & gluons are bound inside
color neutral particles



valence quarks define the hadron
also a **sea** of (anti)quarks & gluons

and what's inside protons and neutrons?

fundamental particles which interact
via the strong force

quarks and gluons

confinement

quarks & gluons are bound inside
color neutral particles



valence quarks define the hadron
also a **sea** of (anti)quarks & gluons

confinement makes the strong force hard to study because the details are locked inside the protons and neutrons

strong force at high temperature

a system that's hot and dense enough for the quarks and gluons to not be confined anymore

strong force at high temperature

a system that's hot and dense enough for the quarks and gluons to not be confined anymore

nucleus
(many protons & neutrons)



strong force at high temperature

a system that's hot and dense enough for the quarks and gluons to not be confined anymore

nucleus
(many protons & neutrons)



+ energy

strong force at high temperature



to create a system that's hot and dense enough for the quarks and gluons to not be confined anymore: the **quark gluon plasma**

Colliders at BNL and CERN

RHIC

0.200 TeV collision energy
Au+Au

LHC



2.76 TeV collision energy
Pb+Pb

relativistic **heavy** ion collisions

Quark Gluon Plasma



relativistic **heavy** ion collisions

Quark Gluon Plasma



lasts for a billionth of a trillionth of a second
and billion times smaller than a pixel on an iPhone display

relativistic **heavy** ion collisions

Quark Gluon Plasma



lasts for a billionth of a trillionth of a second
and billion times smaller than a pixel on an iPhone display

when two individual nuclei collide they create a droplet of this matter
this process happens thousands of times a second

relativistic **heavy** ion collisions

Quark Gluon Plasma



lasts for a billionth of a trillionth of a second
and billion times smaller than a pixel on an iPhone display

when two individual nuclei collide they create a droplet of this matter
this process happens thousands of times a second

we watch the collisions, we cannot do anything external to it

what do we see?

hundreds or thousands of **new** particles are created in each collision

$E = mc^2$



these particles provide the only window into the earlier stages of the collision

we look at each collision individually, but measure billions of collisions!

RHIC @ Brookhaven



PHENIX Detector



PHENIX Detector



PHENIX Detector



the aftermath of a collision



the aftermath of a single collision




STAR

collision geometry



view: one nuclei going into the screen and one coming out







collision geometry



view: one nuclei going into the screen and one coming out

varying the distance between the nuclei, changes the shape and size of the region where the nuclei overlap







collision geometry



view: one nuclei going into the screen and one coming out

varying the distance between the nuclei, changes the shape and size of the region where the nuclei overlap







the parts of the nuclei that don’t overlap continue on and don’t play a role

counting particles





collision geometry





collision geometry





collision geometry





more particles come out the long side than the short side!

interactions are important



interactions are important



liquid rather than a gas



liquid rather than a gas



steep pressure change

liquid rather than a gas



liquid rather than a gas

gradual pressure change



characterizing a liquid

- liquids flow

characterizing a liquid

- liquids flow

  low viscosity

characterizing a liquid

- liquids flow

low viscosity

high viscosity

liquid QGP

liquid QGP

QGP flows well!





liquid QGP

QGP flows well!





low viscosity

$\eta/s > 25\ (1/4\pi)$

liquid QGP

QGP flows well!





low viscosity

$\eta/s > 25\ (1/4\pi)$

$\eta/s(QGP) < 5\ (1/4\pi)$

liquid QGP

QGP flows well!





low viscosity

$\eta/s > 25\ (1/4\pi)$

$\eta/s(QGP) < 5\ (1/4\pi)$

string theory calculation:
universal minimum
$\eta/s > 1/4\pi$

liquid QGP

QGP flows well!





low viscosity

$\eta/s > 25\ (1/4\pi)$

$\eta/s(QGP) < 5\ (1/4\pi)$

string theory calculation:
universal minimum
$\eta/s > 1/4\pi$

determining $\eta/s(QGP)$
is very important

shape changes and particle distributions









ATLAS PLB 707 330 (2012)

isolating shape effects



isolating shape effects



characterizing particle distributions



characterizing particle distributions



v_2 is the strength of the modulation

v_2 in heavy ion collisions



PHENIX PRL 105 062301
CMS PRC 87 014902

quantifying shapes

eccentricity (ε_2) is related to how elongated any shape is



$\varepsilon_2 = 0$ $\varepsilon_2 = 0.17$ $\varepsilon_2 = 0.50$ $\varepsilon_2 = 1$

ratio: v_2 / ε_2



ratio: v_2 / ε_2



relationship between geometry (ε_2) and v_2 is a signature of small viscosity QGP

How Small can the Quark Gluon Plasma Be?

why take something so small and make it smaller?

- changing the shape and size of the QGP help to measure the viscosity



why take something so small and make it smaller?

- is a QGP this small possible?
- could be an extreme variation of the size and time evolution from heavy ion collisions



v2 in p+Pb collisions @ the LHC





ALICE PLB 719 29 (2013)
ATLAS PRL 110 182302 (2013)

big vs small collisions

large ε_2



small ε_2



big vs small collisions

large ε_2



small ε_2



can we have a collision with large eccentricity, but similar size to p+Pb?

varying the small nucleus

d+Au
large ε_2



p+Pb
small ε_2



deuteron (d): 1 proton and 1 neutron

which way does the ellipse go?



in any given event, we can’t control it and it’s hard to measure for these small systems

looking for v2 in d+Au



correlations between pairs of particles
each particle knows something about the collision orientation, but the precision is low

also, there are lots of reasons for particles to be correlated



looking for v2 in d+Au



correlations between pairs of particles
each particle knows something about the collision orientation, but the precision is low

also, there are lots of reasons for particles to be correlated



signal + background

looking for v2 in d+Au



correlations between pairs of particles
each particle knows something about the collision orientation, but the precision is low

also, there are lots of reasons for particles to be correlated



signal + background

background

hunting down the signal

correlations between pairs of particles

each particle knows something about the collision orientation, but the precision is low





hunting down the signal

correlations between pairs of particles

each particle knows something about the collision orientation, but the precision is low





$v_2^2 \cos 2\Delta\phi$

v_2 / ε_2, expectations in d+Au



ALICE PLB 719 29 (2013)
ATLAS PRL 110 182302 (2013)

v_2 / ε_2, expectations in d+Au



ALICE PLB 719 29 (2013)
ATLAS PRL 110 182302 (2013)

v_2 / ε_2



PHENIX PRL 111 212301 (2013)
ALICE PLB 719 29 (2013)
ATLAS PRL 110 182302 (2013)

a small QGP?



continuous behavior from big to small collisions

a small QGP?



continuous behavior from big to small collisions

a small QGP?



continuous behavior from big to small collisions

a small QGP?



continuous behavior from big to small collisions

particle distributions reflect initial shape





in big & small collisions…

each nucleus is a little different

- 200 protons and neutrons move around within the nucleus



each collision is unique



each collision is unique









each collision is unique





shape control

small ε_2





large ε_2



shape control

small ε_2







large ε_2



shape control

small ε_2





large ε_2



small ε_3





shape control

small ε_2





large ε_2



small ε_3





large ε_3



a triangular nucleus?

deuteron: 1 proton, 1 neutron
helium 3 (^{3}He): 2 protons, 1 neutron

a triangular nucleus?

deuteron: 1 proton, 1 neutron
helium 3 (^{3}He): 2 protons, 1 neutron



a triangular nucleus?

deuteron: 1 proton, 1 neutron
helium 3 (^{3}He): 2 protons, 1 neutron



p+Au



d+Au



^{3}He+Au



a triangular nucleus?

deuteron: 1 proton, 1 neutron
helium 3 (^{3}He): 2 protons, 1 neutron



p+Au



d+Au



^{3}He+Au



very successful 3He+Au run concluded last week, analysis in progress!

jets in proton-proton collisions



probing the QGP with jets

p+p collisions



probing the QGP with jets



probing the QGP with jets



probing the QGP with jets



jet quenching



ATLAS PRL105 252303

jet quenching



ATLAS PRL105 252303



jet quenching





ATLAS PRL105 252303

where does the energy not found in the jets end up?
what does that tell us about the matter we're studying?

jet detectors

PHENIX



sPHENIX (coming in 2020)



jet detectors

PHENIX



sPHENIX (coming in 2020)



- **between RHIC & LHC initial QGP temperature is changed**
- **what does the combination of flow and jets tell us about how the QGP works?**

exploring the strong force

- creating a picture of the quark-gluon plasma by using the geometry and variations of the nuclei collided at RHIC and the LHC
  - very small nuclei are providing a unique control of the geometry
  - excited to be able to fully exploit this technique at RHIC with p+Au, d+Au, and 3He+Au collisions soon!



acknowledgements



- all the members of the PHENIX Collaboration
- my colleagues at BNL
- funded through DOE Office of Science

investigating initial state of the nucleus?



investigating initial state of the nucleus?



investigating initial state of the nucleus?



- electrons are point-like particles

investigating initial state of the nucleus?



- electrons are point-like particles



eRHIC

upgrade to allow electrons at RHIC timescale ~ 2025