

Measurement of the W boson mass using 1 fb^{-1} of Dzero data from Run II of the Fermilab Tevatron



Jan Stark

Laboratoire de Physique Subatomique et de Cosmologie
Grenoble, France

Particle Physics Seminar
Brookhaven, December 3, 2009





IN2P3
Les deux infinis

Motivation

W mass is a key parameter in the Standard Model. This model does not predict the value of the W mass, but it predicts this **relation between the W mass and other experimental observables**:

$$M_W = \sqrt{\frac{\pi\alpha}{\sqrt{2}G_F} \frac{1}{\sin\theta_W \sqrt{1-\Delta r}}}$$

Radiative corrections (Δr) depend on M_t as $\sim M_t^2$ and on M_H as $\sim \log M_H$. They include diagrams like these:



Precise measurements of M_W and M_t constrain SM Higgs mass.


For equal contribution to the Higgs mass uncertainty need:

$$\Delta M_W \approx 0.006 \Delta M_t.$$


The limiting factor here will be ΔM_W , not ΔM_t !

Additional contributions to Δr arise in various extensions to the Standard Model,

e.g. in SUSY:



Motivation



For equal contribution to the Higgs mass uncertainty need:

$$\Delta M_W \approx 0.006 \Delta M_t.$$

Current Tevatron average:


$$\Delta M_t = 1.3 \text{ GeV}$$

⇒ would need: $\Delta M_W = 8 \text{ MeV}$

Currently have: $\Delta M_W = 25 \text{ MeV}$

At this point, i.e. after all the precise top mass measurements from the Tevatron, the limiting factor here is ΔM_W , not ΔM_t .



Secret hopes



... as shown by Terry Wyatt at the EPS 2007 conference.

Current precision

Mass of the W Boson




The current world average is still dominated by the final LEP2 results.

The Tevatron average is driven by a recent Run II measurement from CDF (200 pb^{-1}), but the analysis of the Tevatron Run II data is really just starting ...


CDF Run II (200 pb^{-1}):
 $m(W) = 80.413 \pm 0.048 \text{ GeV}$
 Phys.Rev.Lett.99:151801 (2007)
 Phys.Rev.D77:112001 (2008)

Signature in the detector

Isolated, high p_T leptons,
missing transverse momentum in W's




Z events provide excellent
control sample




In a nutshell, measure two objects in the detector:

- Lepton (in principle e or μ ; e in our analysis),
need energy measurement with 0.2 per-mil precision (!!)
- Hadronic recoil, need $\sim 1\%$ precision

Experimental observables



-  No $P_T(W)$
 -  $P_T(W)$ included
 -  Detector Effects added
- $p_T(e)$ most affected by $p_T(W)$




$$M_T = \sqrt{2E_T^l \cancel{E}_T (1 - \cos \Delta\phi)}$$

M_T most affected by measurement of missing transverse momentum

Measurement strategy

W mass is extracted from transverse mass, transverse momentum and transverse missing momentum:

Need Monte Carlo simulation to predict shapes of these observables for given mass hypothesis



NLO event generator : DØ uses **ResBos** [Balazs, Yuan; Phys ReV D56, 5558] + **Photos** [Barbiero, Was; Comp Phys Com 79, 291] for W/Z production and decay

+
Parametrised detector model

↓
W mass templates

+
backgrounds

data → **binned likelihood fit**

↓
W mass

Detector calibration

- calorimeter energy scale
- recoil

First DØ Run II measurement of the W boson mass



1 fb⁻¹ of data
using central electrons ($|\eta| < 1.05$)

~ 500k W events
~ 19k Z events

“blind” analysis : central value hidden but not the uncertainties
Standard blinding technique “à la BaBar”


Unblinding has been done only after collaboration approval



The Tevatron

Proton-antiproton
collisions with
centre-of-mass = 1.96 TeV

36 p and pbar bunches
396 ns between bunch
crossing




Currently the only place in the world where W and Z bosons
can be produced directly

DØ: integrated luminosity


Tevatron is running extremely well
Thanks, FNAL (and Accelerator Division!)

DØ Monthly Data Taking Efficiency 19 April 2002 - 31 October 2009




The DØ detector too !



DØ Run II Integrated Luminosity 19 April 2002 - 30 November 2009



The upgraded Dzero detector



Overview of the calorimeter




46000 cells

50 dead channels

- Liquid argon active medium and (mostly) uranium absorber
- Hermetic with full coverage : $|\eta| < 4.2$
- Segmentation (towers): $\Delta \eta \times \Delta \phi = 0.1 \times 0.1$
(0.05x0.05 in third EM layer, near shower maximum)

Keep in mind: the CAL is not alone !





Samples and weights

The plot on the right shows the average longitudinal profile of a shower with $E = 45$ GeV. Assuming normal incidence, the position of the active parts of the CC are also indicated.

In the reconstruction, we apply artificially high weights to the early layers (especially EM1) in an attempt to partially compensate the losses in the dead material:

Layer	depth (X_0)	weight (a.u.)	weight/ X_0
EM1	2.0	31.199	15.6
EM2	2.0	9.399	4.7
EM3	6.8	25.716	3.8
EM4	9.1	28.033	3.1
FH1	≈ 40	24.885	≈ 0.6




The lower plot illustrates the situation for the same average shower, but this time under a more extreme angle of incidence (physics $\eta = 1$). The shower maximum is now in EM1 !

Energy-dependence and fluctuations

The plots on the previous slide show the *average* shower profile at $E = 45$ GeV.


The plot on the right is basically the same, except that it includes typical *shower fluctuations*.

=> The fraction of energy lost in the dead material varies from shower to shower.



The bottom plot illustrates the situation at a different, lower, energy. The position of the shower maximum (in terms of X_0) varies approximately like $\ln(E)$.


=> The average fraction of energy lost in dead material, as well as the relative importance of shower-by-shower fluctuations depend on the energy of the incident electron.



Average response ...

So we need to apply an **energy-loss correction** to our reconstructed electron energies to account for the energy lost in front of the calorimeter. This correction, as a function of energy and angle (η) is estimated using detailed **detector simulations based on Geant**.

This is the energy correction factor that gets us back to the energy of the incident electron.




This is the energy as reconstructed in the CAL.

Knowing the amount of dead material is the key to energy response linearity: Measure amount of dead material *in situ* using electrons from $Z \rightarrow e e$.

... and fluctuations around the average


Here we show the impact on the energy resolution for electrons. This is again from a detailed detector simulation based on Geant.

Resolution at normal incidence, as a function of electron energy:



for an ideal sampling calorimeter (no dead material) one would expect this to scale as $1/\sqrt{E}$

Resolution at $E = 45$ GeV, as a function of the angle of incidence (η):



for an ideal sampling calorimeter (no dead material) one would expect this to be almost flat

How to split our (already small) $Z \rightarrow e e$ sample ??

So we need to understand both average response and the resolution as a function of both energy and angle of incidence.

$Z \rightarrow e e$ data gives us access to a line in energy/angle space. Consider CC/CC events. At a given angle, the distribution of energies provided by Nature is rather narrow.


How to proceed:

- => Bin electrons in angle (5 bins).
- => Two electrons per Z.
- => 15 distinct combinations of bins - “categories” (no E ordering).

Split **CC/CC** $Z \rightarrow e e$ sample into the 15 categories and study measured **Z mass and mass resolution per category**.

Once the information from Z has been harvested, we still need to **propagate that down to the lower energies of the W**.

Need to understand scaling laws.




bin 0 : $0 \leq \eta < 0.2$
bin 1 : $0.2 \leq \eta < 0.4$
bin 2 : $0.4 \leq \eta < 0.6$
bin 3 : $0.6 \leq \eta < 0.8$
bin 4 : $0.8 \leq \eta $

Category	Bins of Each Electron
10	0-0
11	0-1
12	0-2
13	0-3
14	0-4
15	1-1
16	1-2
17	1-3
18	1-4
19	2-2
20	2-3
21	2-4
22	3-3
23	3-4
24	4-4

Simple plots (after splitting)

Let's start with a few simple plots that are based on the idea of splitting the sample according to eta of the two electrons. Here are the **Z mass peaks (early version of data reconstruction)** for “**both electrons very central**” and “**both electrons very forward**”, i.e. “**both electrons at close to normal incidence**” and “**both electrons at highly non-normal incidence**”



We note:

- different resolutions (material !),
- the peaks are not in the same place.


Why aren't the peaks in the same place ? Could be a problem in the MC-based E-loss corrections. But could also be a problem with gain calibrations in different regions of the CAL. This plot alone is not going to tell us, we need more information, new observables.

Need more information: additional observables

Let's go back to one of the plots that we have discussed on an earlier slide.

It clearly suggests that we should try to **exploit the longitudinal segmentation of the EM CAL** to get a handle on dead material:

Imagine we vary the size of the “DEAD” region a little bit
=> the individual layers (EM1 etc) would sample different parts of the shower and therefore see different fractions of the shower energy !!




Using the longitudinal segmentation to get a handle on material is a standard technique, it is discussed in the textbooks (e.g. Wigmans).

Back to Dzero. Let's compare data (old reconstruction) and full Monte Carlo (nominal geometry) in terms of the four fractional EM energy deposits. We do this separately in each of the 15 eta categories.

DO Before tuning of material model

Before tuning of material model:
distributions of fractional energy deposits
do not quite match between data and the simulation.




Fit for amount of missing material

“Turn the plots from the previous slides into a fit for the amount of missing material”:

Take data/MC ratios per η category for EM1, EM2 and EM3 and fit each one (separately) to a constant. Add the chi-squareds from the three fits. Vary amount of extra material to minimise the global chi-squared.

This implies that we leave the absolute energy scale of each layer free to float. This is because this fit is the first time that we have a handle on the intercalibration of the layers.




Amount of fudge material to within less than $0.01X_0$!

With comparatively small systematics from background (underlying event) subtraction and modelling of cut efficiencies.




After tuning of material model

After tuning of material model:
distributions of fractional energy deposits
are very well described by the simulation.




As a cross-check:
Repeat fit for nX_0 , separately for
each EM layer. Good consistency
is found.



Correction to the raw energy

An **energy-loss correction** is applied to our reconstructed electron energies to account for the energy lost in front of the calorimeter. This correction, as a function of energy and angle (η) is estimated using detailed **detector simulations based on Geant** including the fitted amount of missing material.

This is the energy correction factor that gets us back to the energy of the incident electron.



This is the energy as reconstructed in the CAL.

This energy correction is applied on the data and not parameterised in our fast MC.



Electrons: energy scale

After having corrected for the effects of the uninstrumented material:
final energy response calibration, using $Z \rightarrow e e$, the known Z mass value from LEP,
and the standard “ f_z method”:

$$E_{\text{measured}} = \alpha \times E_{\text{true}} + \beta$$

Use energy spread of electrons in Z decay to constrain α and β .

In a nutshell: the f_z observable allows you to split your sample of electrons from $Z \rightarrow e e$ into subsamples of different true energy; this way you can “scan” the electron energy response as a function of energy.

$$f_z = (E(e1) + E(e2))(1 - \cos(\gamma_{ee})) / m_Z$$

γ_{ee} is the opening angle between the two electrons

Result:

$$\begin{aligned} \alpha &= 1.0111 \pm 0.0043 \\ \beta &= -0.404 \pm 0.209 \text{ GeV} \\ \text{correlation: } &-0.997 \end{aligned}$$

This corresponds to the dominant systematic uncertainty (by far) in the W mass measurement (but this is really just Z statistics ... more data will reduce it) :

$$\Delta m(W) = 34 \text{ MeV, } 100 \% \text{ correlated between all three observables}$$



Electrons: energy resolution

Electron energy resolution is driven by two components:
sampling fluctuations and constant term


Sampling fluctuations are driven by sampling fraction of CAL modules (well known from simulation and testbeam) and by uninstrumented material. As discussed before, amount of material has been quantified with good precision.

Constant term is extracted from $Z \rightarrow e e$ data (essentially fit to observed width of Z peak).


Result:

$$C = (2.05 \pm 0.10) \%$$

in excellent agreement with Run II design goal (2%)



Switching gears: recoil model



Recoil model

Recoil vector in parameterised MC: $\vec{u}_T = \vec{u}_T^{\text{Hard}} + \vec{u}_T^{\text{Soft}} + \vec{u}_T^{\text{Elec}} + \vec{u}_T^{\text{FSR}}$

$$\vec{u}_T^{\text{Hard}} = \vec{f}(\vec{q}_T)$$

Hard component that balances the vector boson in transverse plane.

Ansatz from full $Z \rightarrow \nu \nu$ MC; plus free parameters for fine tuning, e.g. multiplicative scale adjustment as function of q_T :

$$\text{RelResp} = \text{RelScale} + \text{RelOffset} \cdot \exp \frac{-q_T}{\tau_{\text{HAD}}}$$

$$\vec{u}_T^{\text{Soft}} = \alpha_{\text{MB}} \cdot \vec{E}_T^{\text{MB}} + \alpha_{\text{ZB}} \cdot \vec{E}_T^{\text{ZB}}$$

Soft component, not correlated with vector boson.

Two sub-components; - additional ppbar interactions and detector noise: from ZB events, plus parameter for fine tuning
- spectator partons: from MB events, plus parameter for fine tuning

$$\vec{u}_T^{\text{Elec}} = - \sum_e \Delta u_{\parallel} \cdot \hat{p}_T(e)$$

Recoil energy “lost” into the **electron cones**.
Electron energy leakage outside cluster.





$$\vec{u}_T^{\text{FSR}} = \sum_{\gamma} \vec{p}_T(\gamma)$$

FSR photons (internal bremsstrahlung) outside cone; includes detailed response model.



Recoil calibration


Final adjustment of free parameters in the recoil model is done *in situ* using balancing in $Z \rightarrow e e$ events and the standard UA2 observables.



UA2 observables:
In transverse plane, use a coordinate system defined by the bisector of the two electron momenta.

$$\eta\text{-imbalance} : (\vec{P}_t^{ee} + \vec{P}_t^{rec}) \cdot \hat{\eta}$$

$$\xi\text{-imbalance} : (\vec{P}_t^{ee} + \vec{P}_t^{rec}) \cdot \hat{\xi}$$







Examples: η_{imb} distributions

$1 < p_T(ee) < 2 \text{ GeV}$

$20 \text{ GeV} < p_T(ee)$







Electron reco efficiency model

Efficiency model also takes into account **relative orientation** of electron and “rest of the event” (hadronic activity). For example:

- Efficiency corrections vs. $p_T(e)$ and scalar E_T .
- Efficiency corrections vs. $u_{||}$.


Much of this level of detail is only necessary for a measurement of the W width, not the mass.






Results: $Z \rightarrow e e$ data


ZCandMass_CCCC_Trks




ZCandElecPt_0



ZCandPt_0







ZCandRecoilPt_0



✓ Good agreement between parameterised MC and collider data.



Mass fits







$$m(Z) = 91.185 \pm 0.033 \text{ GeV (stat)}$$

(remember that Z mass value from LEP was an input to electron energy scale calibration, PDG: $m(Z) = 91.1876 \pm 0.0021 \text{ GeV}$)

$$m(W) = 80.401 \pm 0.023 \text{ GeV (stat)}$$



Mass fits



$$m(W) = 80.400 \pm 0.027 \text{ GeV (stat)}$$

$$m(W) = 80.402 \pm 0.023 \text{ GeV (stat)}$$




Summary of uncertainties

systematic uncertainties

Source	$\sigma(m_W)$ MeV m_T	$\sigma(m_W)$ MeV p_T^e	$\sigma(m_W)$ MeV \cancel{E}_T
Experimental			
Electron Energy Scale	34	34	34
Electron Energy Resolution Model	2	2	3
Electron Energy Nonlinearity	4	6	7
W and Z Electron energy loss differences (material)	4	4	4
Recoil Model	6	12	20
Electron Efficiencies	5	6	5
Backgrounds	2	5	4
Experimental Total	35	37	41
W production and decay model			
PDF	9	11	14
QED	7	7	9
Boson p_T	2	5	2
W model Total	12	14	17
Total	37	40	44
statistical	23	27	23
total	44	48	50

Comparison to previous results




The new result from DØ is the **single most precise measurement** of the W boson mass to date.

On the left we quote our m_T result. A combination with the (correlated !) results from the other two observables reduces the uncertainty from 44 MeV to 43 MeV.

The new result is in good agreement with previous measurements.

Latest constraints on Higgs mass


For detailed description of this update, please see web page of Gfitter group.




Back to Terry's hopes

... as shown at the EPS 2007 conference.

Are such expectations reasonable ?



Projection



**CDF/DØ combined
 10 fb⁻¹ 2011 (?)**

$\Delta m_W \sim 10$ MeV
 $\Delta m_{top} \sim 1$ GeV



Summary and outlook

We have presented a **new measurement of the W boson mass** from the DØ Collaboration. It is based on central electrons in 1 fb^{-1} of Run II data:

$$\begin{aligned} m_W &= 80.401 \pm 0.023(\text{stat}) \pm 0.037(\text{syst}) \text{ GeV} = 80.401 \pm 0.044 \text{ GeV} \quad (m_T) \\ &80.400 \pm 0.027(\text{stat}) \pm 0.040(\text{syst}) \text{ GeV} = 80.400 \pm 0.048 \text{ GeV} \quad (p_T^e), \\ &80.402 \pm 0.023(\text{stat}) \pm 0.044(\text{syst}) \text{ GeV} = 80.402 \pm 0.050 \text{ GeV} \quad (\cancel{E}_T). \end{aligned}$$



A combination of the results from the three observables gives:

$$\mathbf{80.401 \pm 0.021 \text{ (stat.)} \pm 0.038 \text{ (syst.)} = 80.401 \pm 0.043 \text{ GeV}}$$

This is the **most precise single measurement** of the W boson mass to date.

This measurement is in **good agreement** with a previous Run II measurement from CDF (electron and muons in 200 pb^{-1} of data), as well as with the LEP average.

DØ and CDF use **very different techniques** for the main ingredient of the measurement, namely to establish the lepton energy scale. Their systematic uncertainties are **uncorrelated** to a large extent, which is good for **cross-checks and combination**. Similar comments apply to (non-)correlation with LEP results.

For both DØ and CDF these measurements are just the beginning. Both collaborations are analysing larger datasets. CDF predict 25 MeV total uncertainty with 2.3 fb^{-1} . DØ expect similar or better uncertainties with the additional 5 fb^{-1} in the can.

Backup slides

Comments on analysis strategy

Before analysing the collider data, we perform a **Monte Carlo closure test**. This means we treat simulated events from a detailed Pythia/Geant simulations as collider data and perform a full W mass/width analysis. Goal: develop and test analysis procedures and code with known input values. At each analysis step, check that predictions from parameterised MC match MC truth.

We perform our measurements as a **blind analysis**. This means that the central values (but not the uncertainties) are deliberately hidden from the analysers and reviewers until the analysis is considered complete. The blinding technique we used is a standard technique that is routinely used by other collaborations, *e.g.* BaBar:




Simply change your mass fitting program in such a way that it reports the fitted mass, offset by some hidden offset.

The offset is the same for all three observables (\Rightarrow allow comparisons), no uncertainties, neither statistical nor systematic are ever obscured by the blinding.

“Unblinding” has been done only after collaboration approval.


MC closure test: $Z \rightarrow e e$

✓ Good agreement between full and parameterised MC.




MC closure test: $W \rightarrow e \nu$


WCandMt_Spatial_Match_0




WCandElecPt_Spatial_Match_0




WCandMet_Spatial_Match_0




χ distribution with overall $\chi^2 = 89.8$ for 100 bins




χ distribution with overall $\chi^2 = 79.2$ for 70 bins



χ distribution with overall $\chi^2 = 85.0$ for 70 bins



MC closure test: $W \rightarrow e \nu$



Model of W production and decay

Tool	Process	QCD	EW
RESBOS	W, Z	NLO	-
WGRAD	W	LO	complete $\mathcal{O}(\alpha)$, Matrix Element, ≤ 1 photon
ZGRAD	Z	LO	complete $\mathcal{O}(\alpha)$, Matrix Element, ≤ 1 photon
PHOTOS			QED FSR, ≤ 2 photons


Our main generator is “ResBos+Photos”. The NLO QCD in ResBos allows us to get a reasonable description of the p_T of the vector bosons. The two leading EWK effects are the first FSR photon and the second FSR photon. Photos gives us a reasonable model for both.

We use W/ZGRAD to get a feeling for the effect of the full EWK corrections.

The final “QED” uncertainty we quote is 7/7/9 MeV (m_T, p_T, MET).


This is the sum of different effects; the two main ones are:

- Effect of full EWK corrections, from comparison of W/ZGRAD in “FSR only” and in “full EWK” modes (5/5/5 MeV).
- Very simple estimate of “quality of FSR model”, from comparison of W/ZGRAD in FSR-only mode vs Photos (5/5/5 MeV).




This is a U/LAr sampling calorimeter

More detailed view of one CC-EM module :



incident particle

sampling fraction: 15 %




Basically a stack of Uranium plates with liquid Argon in between. Shower develops in U and LAr (mainly U); charged shower particles ionise the Argon atoms => current in Argon because of HV applied across each gap. This current is measurable (thanks to electronic charge amplifiers with very large gain).


EM1, EM2, EM3 and EM4 are read out separately; each one of these layers regroups a number of digaps.

Basics of the readout

- Detector signal ~ 450 ns long
(bunch crossing time: 396 ns)
- Charge preamplifiers
- BLS (baseline subtraction) boards
 - short shaping of $\sim 2/3$ of integrated signal
 - signal sampled and stored every 132 ns in analog buffers (SCA) waiting for L1 trigger
 - samples retrieved on L1 accept, then baseline subtraction to remove pile-up and low frequency noise
 - signal retrieved after L2 accept
- Digitisation



Have ability to sample and record the shaped signal also at (320 ± 120) ns to make sure we are on the peak.




$$J/\Psi \rightarrow e^+ e^-$$

Fortunately, when I said “*extrapolation*” down to the W , that was not the whole story. We also have another di-electron resonance that sits **lower** in energy than the W : the J/Ψ .

At a hadron collider, such a sample is *extremely* hard to obtain. One of the keys to our success is D0's excellent *Central Track Trigger*. It allows us to trigger on isolated tracks already at Level 1. We typically require two tracks of $p_T > 3$ GeV.

It took us many many person-months to obtain this sample: design/implementation of the trigger, understanding efficiencies, etc, etc.

JPsi Resonance for LOW Triggers (Entire CC)




In contrast to the Z , the energy resolution at J/Ψ energies is practically insensitive to issues with gain calibration (the constant term in the energy resolution is irrelevant). The J/Ψ is a nice probe for sampling fluctuations and scale issues related to dead material.

Cross-sections: QED is easy, right ?


But in practice these calculations involve time-consuming Hartree-Fock calculations, partial wave expansions, etc, etc.

In addition, popular simulation programs (like Geant or EGS) often use simplified models or simple parameterisations of cross sections in order to avoid large look-up tables and to implement fast random number techniques.

A detailed comparison of Geant and EGS to state-of-the-art cross section calculations is striking, especially for Bremsstrahlung.




Example: Bremsstrahlung in by electrons in uranium



T = kinetic energy of incident electron k = energy of the radiated photon


EM fractions in $Z \rightarrow e^+ e^-$ events

Use electrons from $Z \rightarrow e e$, plot mean fractional energy deposit in each one of the EM layers. Separate the events into the standard categories in physics eta. The plot below shows each of the four EM fractions for each of the 15 categories.



This is a busy plot that can be tricky to read. Let's look at the data/MC ratios instead (on the next slide).


EM fractions in $Z \rightarrow e^+ e^-$ events



Clear trends are visible, especially for EM1 and EM3.

Also, the excursions away from unity are pretty large. Part of the mean per-layer excursion could be explained by the layers not being properly calibrated with respect to each other, but deviations of O(5 %) are not really expected.

EM fractions in $Z \rightarrow e^+ e^-$ events



Certainly less trendy than with the nominal detector geometry.

The layers that receive the bulk of the energy (EM1, EM2 and EM3) are also much closer to unity.

Dzero Run I


DØ Collaboration, PRD 58, 092003 (1998)

Observable: “transverse mass”

$$M_T = \sqrt{2E_T^l E_T (1 - \cos \Delta\phi)}$$

Relatively robust against uncertainties in physics model.

Uncertainties



Model detector effects using parameters “from data” (and a lot of hypotheses).

Generate M_T templates for different M_T points \rightarrow likelihood fit.


Understanding the detector behaviour, based mainly on $Z \rightarrow e e$ and $\Psi \rightarrow e e$ calibration samples, is crucial.

Stat.	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
W sample	70	85	105
Z sample	65	65	65
Total	95	105	125

“Detector understanding”	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
Calorimeter linearity	20	20	20
Calorimeter uniformity	10	10	10
Electron resolution	25	15	30
Electron angle calibration	30	30	30
Electron removal	15	15	20
Selection bias	5	10	20
Recoil resolution	25	10	90
Recoil response	20	15	45
Total	60	50	115

“Production and decay model”	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
$p_T(W)$ spectrum	10	50	25
Parton distribution functions	20	50	30
Parton luminosity β	10	10	10
Radiative decays	15	15	15
W width	10	10	10
Total	30	75	45

$W \rightarrow e \nu$ data WARNING !




Data-MC apparently not in good agreement !

Error bars only reflect limited W statistics they do not reflect the much more limited **Z statistics** that have been used to calibrate the recoil model !


The blue band in the bottom plot reflects one sigma excursions in the recoil parameters.

→ agreement is not so bad !




$Z \rightarrow e e$ and $W \rightarrow e \nu$

Data in red
MC in blue




SET

GeV




GeV



u_T


GeV





GeV

$Z \rightarrow e e$ and $W \rightarrow e \nu$


Data in red
MC in blue



u_{para}





u_{perp}




Stability checks

Changes in the fitted m_W when the fitting range (m_T observable) is varied.




Stability checks

Instantaneous luminosity (split data into two subsets – high and low inst. luminosity)






Time (i.e. data-taking period)






Sorry, plots still in terms of blinded mass, but it does not matter here.

Stability checks

Scalar E_T (“global event activity as seen by calorimeter”)




Electron distance from phi cracks



Sorry, plots still in terms of blinded mass, but it does not matter here.


Stability checks

Cut on u_T (“length of recoil vector”)



Sorry, plots still in terms of blinded mass, but it does not matter here.

The new world average



Latests constraints on Higgs mass

For detailed description of this update, please see web page of Gfitter group.

